

On the KP Hierarchy, $\hat{W}_\infty$ Algebra, and Conformal $SL(2, \mathbb{R})/U(1)$ Model — The Classical and Quantum Cases¹

Feng Yu[†] and Yong-Shi Wu^{††}

[†] *Institute for Theoretical Physics, State University of New York
Stony Brook, New York 11794, U.S.A.*

^{††} *Center for Theoretical Physics, MIT
Cambridge, MA 02139, U.S.A.*

and

*Department of Physics, University of Utah
Salt Lake City, Utah 84112, U.S.A.*

ABSTRACT

We give a unified description of our recent results on the inter-relationship between the integrable infinite KP hierarchy, nonlinear $\hat{W}_\infty$ current algebra and conformal noncompact $SL(2, \mathbb{R})/U(1)$ coset model both at the classical and quantum levels. In particular, we present the construction of a quantum version of the KP hierarchy by deforming the second KP Hamiltonian structure through quantizing the $SL(2, \mathbb{R})_k/U(1)$ model and constructing an infinite set of commuting quantum $\hat{W}_\infty$ charges (at least at $k=1$).

¹Invited talk given by the second author at the XXI International Conference on Differential Geometric Methods in Theoretical Physics, Nankai Institute of Mathematics, Tianjin, China; June 5-9, 1992; to appear in Proceedings.

Introduction and Summary

One of the most exciting recent developments in mathematical physics is the revelation of the close relationship between infinite dimensional (extended) conformal algebras, nonlinear integrable differential systems and 2d exactly solvable field theories, including conformal field models, 2d quantum gravity and string theory. A well-known example is the Virasoro algebra, which is known to be isomorphic to the second Hamiltonian structure of the KdV hierarchy, and appears as the operator product algebra in 2d minimal conformal models, and as constraint algebra in $c < 1$ string theory. Similar relations happen to Zamolochikov's extended W_N algebras ($N \geq 3$), extension of the Virasoro algebra generated by currents of higher spin ($2 \leq s \leq N$).

In our recent study¹⁻⁴, we generalize this profound relationship to the W -infinity algebras, which are generated by currents of all integer spins ($s \geq 1$ or $s \geq 2$) including the energy-momentum tensor and can be viewed as (non-unique) large N limits of the W_N algebras. There are some very deep connection between such algebras and the famous KP hierarchy, which is known to contain all generalized KdV hierarchies, and the conformal $SL(2, R)/U(1)$ coset model that supports the black-hole solution in $D = 2$ string theory.

The results reported in our previous papers¹⁻⁴ can be summarized as follows: 1) The centerless linear $W_{1+\infty}$ algebra can be identified (also see ref. 5) as the first Hamiltonian structure of the KP hierarchy. 2) There exists a unique non-linear deformation² of the W_∞ algebra under certain homogeneity assumptions, which we call $\hat{W}_\infty$ and is believed to contain all W_N by reduction. 3) This $\hat{W}_\infty$ algebra is identified as the (modified) second KP Hamiltonian structure (see also ref. 6). 4) We have found a two boson realization of the nonlinear $\hat{W}_\infty$ algebra, in closed form for all higher spin currents. 5) Using it, we discovered a field theoretical realization of $\hat{W}_\infty$ as a hidden current algebra in the 2d $SL(2, R)/U(1)$ coset model. 6) The well-known infinite set of KP Hamiltonians, considered as involutive $\hat{W}_\infty$ charges, generate infinitely many KP flows in the coset model, all of which preserve the $\hat{W}_\infty$ current algebra.

The above results 3)-6) hold true not only at the classical but also the *quantum* level: 7) A consistent quantum deformation of the $\hat{W}_\infty$ algebra, together with its two boson realization, is obtained by quantizing the $SL(2, R)_k/U(1)$ coset model. 8) We

have been able to construct an infinite set of commuting quantum $\hat{W}_\infty$ charges at level $k=1$. 9) These quantum charges are used to generate a quantum KP hierarchy, whose Hamiltonian structure is just the quantum $\hat{W}_\infty$.

In this talk we give a unified description of these results. Two long papers containing detailed proofs will be published soon^{7,8}.

KP Bi-Hamiltonian Structure, $\hat{W}_{1+\infty}$ and $\hat{W}_\infty$

The integrable KP-Sato hierarchy is an infinite set of nonlinear partial-differential equations, which reads in the Lax form

$$\frac{\partial L}{\partial t_m} = [(L^m)_+, L] \quad (1)$$

where L is a first-order pseudo-differential operator (PsD) $L = D + \sum_{r=-1}^{\infty} u_r D^{-r-1}$, with coefficients u_r functions of z and various times $t_m (m = 1, 2, 3, \dots)$, and $D \equiv \partial/\partial z$.

The integrability of the KP hierarchy is a direct consequence of its bi-Hamiltonian structure. To put the KP hierarchy into the Hamiltonian form

$$\frac{\partial u_r}{\partial t_m} = \{u_r(z), \oint H_m(z) dz\}, \quad (2)$$

one needs to define both the Hamiltonian functions $H_m(z)$ and the Poisson brackets

$$\{u_r(z), u_s(z')\} = k_{rs}(u_p(z)) \delta(z - z'). \quad (3)$$

To obtain the bi-Hamiltonian structure, we substitute an arbitrary PsD of order N , $P = \sum_{s=-\infty}^N D^s p_s$, (N can be arbitrarily large) into

$$\hat{K}(P) \equiv (\hat{L}P)_+ \hat{L} - \hat{L}(P\hat{L})_+ = \sum_{r,s=-1}^{\infty} \hat{k}_{rs} p_s D^{-r-1}, \quad (4)$$

where $\hat{L} = L + c$ (c is a real parameter).

Proposition 1: For fixed pair (r, s) , the operator $\hat{k}_{rs}$ becomes stable (i.e. independent of N) when N is sufficiently large. Explicitly, it is given by (for $r, s \geq 0$)

$$\hat{k}_{rs} = \sum_{l=0}^{s+1} \binom{s+1}{l} D^l \hat{u}_{r+s+1-l} - \sum_{l=0}^{r+1} \binom{r+1}{l} \hat{u}_{r+s+1-l} (-D)^l$$

$$\begin{aligned}
& + \sum_{l=0}^{\infty} \left[\sum_{t=l}^{l+s} (-1)^l \binom{t-s-1}{l} \right. \\
& \quad \left. - \sum_{t=r+1}^{r+s+1} \sum_{k=0}^{t-r-1} (-1)^{l+k} \binom{t-k-1}{l-k} \binom{s}{k} \right] \hat{u}_{t-l-1} D^l \hat{u}_{r+s-t}, \\
\hat{k}_{r-1} &= - \sum_{l=1}^r \binom{r}{l} \hat{u}_{r-l} (-D)^l, \\
\hat{k}_{-1s} &= \sum_{l=1}^s \binom{s}{l} D^l \hat{u}_{s-l}, \quad \hat{k}_{-1-1} = -D;
\end{aligned} \tag{5}$$

with $\hat{u}_{-1} = u_{-1} + c$, $\hat{u}_r = u_r$ ($r \geq 0$). And the operator matrix k_{rs} defines a one-parameter family of Hamiltonian structures for functions u_r .

Proposition 2: The following limits

$$\{u_r(z), u_s(z')\}_1 = \lim_{c \rightarrow \infty} \left(\frac{1}{c} \hat{k}_{rs}(z) \right) \delta(z - z') \quad r, s \geq 0; \tag{6}$$

$$\{u_r(z), u_s(z')\}_2 = \lim_{c \rightarrow 0} \hat{k}_{rs}(z) \delta(z - z') \quad r, s \geq -1 \tag{7}$$

coincide, respectively, with the first and second Hamiltonian structures of the KP hierarchy with the Hamiltonian functions

$$H_m^{(1)} = \frac{1}{m+1} \text{Res} L^{m+1}, \quad H_m^{(2)} = \frac{1}{m} \text{Res} L^m. \tag{8}$$

Note the first KP Hamiltonian structure is linear in u_r ($r \geq 0$):

$$\lim_{c \rightarrow \infty} \left(\frac{1}{c} \hat{k}_{rs} \right) = \sum_{l=0}^s \binom{s}{l} D^l u_{r+s-l} - \sum_{l=0}^r \binom{r}{l} u_{r+s-l} (-D)^l. \tag{9}$$

(The u_{-1} current becomes decoupled from the algebra.) Previously we have identified it with the $W_{1+\infty}$ algebra. We call the nonlinear second Hamiltonian structure (7) (with $c = 0$) as the $\hat{W}_{1+\infty}$ algebra, in which the current u_r is considered to have spin $r + 2$ ($r \geq -1$). It is a universal W -algebra, containing all known W -algebras of the infinite type or of the finite type either by contraction or by reduction. However, u_{-1} does not evolve at all: $\partial u_{-1} / \partial t_m = 0$. One can always impose the constraint $u_{-1} = 0$ which is second-class in $\hat{W}_{1+\infty}$.

Proposition 3: With $u_{-1} = 0$, the modified second KP Hamiltonian structure is

$$k_{rs}^{(2D)} = \sum_{l=0}^{s+1} \binom{s+1}{l} D^l u_{r+s+1-l} - \sum_{l=0}^{r+1} \binom{r+1}{l} u_{r+s+1-l} (-D)^l$$

$$\begin{aligned}
& + \sum_{l=0}^{\infty} \left[\sum_{t=l+1}^{l+s} (-1)^l \binom{t-s-1}{l} \right. \\
& - \sum_{t=r+1}^{r+s} \sum_{k=0}^{t-r-1} (-1)^{l+k} \binom{t-k-1}{l-k} \binom{s}{k} \left. \right] u_{t-l-1} D^l u_{r+s-t} \\
& - \sum_{l=1}^r \sum_{k=1}^s (-1)^l \binom{r}{l} \binom{s}{k} u_{r-l} D^{l+k-1} u_{s-k}.
\end{aligned} \tag{10}$$

It can be viewed as the unique nonlinear, centerless deformation of W_{∞} under certain natural homogeneity requirements. We call it as the $\hat{W}_{\infty}$ algebra.

A direct consequence is the following interesting relation between $\hat{W}_{\infty}$ and $W_{1+\infty}$:

$$\oint \{ \hat{W}_{r+1}(z), \hat{W}_s(z') \}_{2D} dz' = \oint \{ W_r(z), W_s(z') \}_1 dz'. \tag{11}$$

Free Field Realization of $\hat{W}_{\infty}$ and $SL(2, R)/U(1)$

The known realization of $W_{1+\infty}$ with a complex fermion is simply summarized by

$$L \equiv D + \sum_{r=0}^{\infty} u_r D^{-r-1} = D - \bar{\psi} D^{-1} \psi. \tag{12}$$

Considering only the spin $s \geq 2$ currents, for the W_{∞} generators we have

$$(LD)_{-} \equiv \sum_{r=0}^{\infty} v_r D^{-r-1} = -(\bar{\psi} D^{-1} \psi D)_{-}. \tag{13}$$

Meanwhile, W_{∞} can be realized in terms of a complex boson

$$L \equiv D + \sum_{r=0}^{\infty} v_r D^{-r-1} = D + \bar{j} D^{-1} j. \tag{14}$$

where the bosonic currents $\bar{j}(z) = \bar{\phi}'(z)$, $j(z) = \phi'(z)$ satisfy

$$\{ \bar{j}(z), j(z') \} = \partial_z \delta(z - z'); \quad \{ \bar{j}(z), \bar{j}(z') \} = \{ j(z), j(z') \} = 0. \tag{15}$$

A nonlinear deformation of this leads to

Proposition 4: The following gives a realization of all currents u_r of $\hat{W}_{\infty}$:

$$\begin{aligned}
L & \equiv D + \sum_{r=0}^{\infty} u_r D^{-r-1} = D + \bar{j} \frac{1}{D - (\bar{j} + j)} j \\
& = D + \bar{j} D^{-1} j + \bar{j} D^{-1} (\bar{j} + j) D^{-1} j + \dots.
\end{aligned} \tag{16}$$

This realization of $\hat{W}_\infty$ naturally emerges in the $SL(2, R)/U(1)$ coset model. Recall the bosonized $SL(2, R)_k/U(1)$ parafermion currents: (classically we set $k = 1$)

$$\psi_+ = \bar{j}e^{\bar{\phi}+\phi}, \quad \psi_- = je^{-\bar{\phi}-\phi}. \quad (17)$$

We propose to expand the bilocal product $\psi_+(z)\psi_-(z')$ in powers of $z - z'$ to all orders.

Proposition 5: The product $\psi_+(z)\psi_-(z')$ generate all two-boson $\hat{W}_\infty$ currents (16):

$$\bar{j}e^{\bar{\phi}+\phi}(z)je^{-\bar{\phi}-\phi}(z') = \sum_{r=0}^{\infty} u_r(z) \frac{(z - z')^r}{r!}. \quad (18)$$

Formally this can also be rewritten as $u_r(z) = \psi_+(z)(-\partial_z)^r\psi_-(z)$, or

$$L = D + \sum_{r=0}^{\infty} u_r D^{-r-1} = D + \psi_+ D^{-1} \psi_-. \quad (19)$$

It follows that the Poisson brackets for $\hat{W}_\infty$ can be generated from

$$\begin{aligned} & \{\bar{j}e^{\bar{\phi}+\phi}(z)je^{-\bar{\phi}-\phi}(z - \epsilon), \bar{j}e^{\bar{\phi}+\phi}(w)je^{-\bar{\phi}-\phi}(w - \sigma)\} \\ \equiv & \{\psi_+(z)\psi_-(z - \epsilon), \psi_+(w)\psi_-(w - \sigma)\} = \sum_{r,s=0}^{\infty} k_{rs}(z)\delta(z - w) \frac{\epsilon^r \sigma^s}{r!s!}. \end{aligned} \quad (20)$$

Involutive KP charges and $\hat{W}_\infty$ Symmetry in $SL(2, R)/U(1)$

The KP charges, $Q_m \equiv \oint H_m(z)dz$, are known to be conserved under the KP flows:

$$\frac{\partial}{\partial t_n} \oint H_m(z, t_k) dz = 0. \quad (21)$$

and they are in involution:

$$\{Q_n^{(2)}, Q_m^{(2)}\}_{2D} = \{Q_{n-1}^{(1)}, Q_m^{(1)}\}_1 = 0. \quad (22)$$

This implies that in the $SL(2, R)/U(1)$ coset model there exist infinitely many commuting $\hat{W}_\infty$ charges, which generate an infinite symmetry

$$\frac{\partial u_r}{\partial t_m} = \{u_r, Q_m\}, \quad \text{or} \quad \delta_m u_r = \epsilon_m \{u_r, Q_m\}. \quad (23)$$

An important property of these symmetry transformations is that the $\hat{W}_\infty$ current algebra for the u_r 's is preserved under these flows:

$$\frac{\partial}{\partial t_m} (\{u_r(z), u_s(z')\} - k_{rs}(z)\delta(z - z')) = 0. \quad (24)$$

Also these symmetries are abelian to each other, in the sense that the KP flow in one charge direction is invariant under the transformation generated by another $\hat{W}_\infty$ charge:

$$\delta_m\left(\frac{\partial u_r}{\partial t_n} - \{u_r, Q_n\}\right) = \epsilon_m\{u_r, \{Q_n, Q_m\}\} = 0. \quad (25)$$

Note that these results hold for the $W_{1+\infty}$ charges as well.

The free boson realization of $\hat{W}_\infty$ allows us to construct the KP flows of j and $\bar{j}$:

$$\frac{\partial \bar{j}}{\partial t_m} = \{\bar{j}, Q_m\}, \quad \frac{\partial j}{\partial t_m} = \{j, Q_m\} \quad (26)$$

(See also ref. 9.) Note that the Hamiltonian structure of this hierarchy is simply the fundamental Poisson brackets (15) for $\bar{j}$ and j , which are invariant under the $\bar{j}$ - j flows:

$$\frac{\partial \{\bar{j}, j\}}{\partial t_m} = \{\{\bar{j}, Q_m\}, j\} + \{\bar{j}, \{j, Q_m\}\} = \{\{\bar{j}, j\}, Q_m\} = 0. \quad (27)$$

Similarly, the free fermion realization (12) of $W_{1+\infty}$ gives the KP Hamiltonians

$$H_1 = \bar{\psi}\psi, \quad H_2 = \frac{1}{2}(\bar{\psi}'\psi - \bar{\psi}\psi'), \quad H_3 = \frac{1}{3}(\bar{\psi}''\psi - \bar{\psi}'\psi' + \bar{\psi}\psi''), \quad \text{etc.} \quad (28)$$

Their charges can also be used to generate a compatible and integrable $\bar{\psi}$ - ψ hierarchy

$$\frac{\partial \bar{\psi}}{\partial t_m} = \{\bar{\psi}, Q_m\}, \quad \frac{\partial \psi}{\partial t_m} = \{\psi, Q_m\}. \quad (29)$$

Quantum $\hat{W}_\infty$ and Its Two Boson Realization

Because of the connection of the $SL(2, R)/U(1)$ model to the black holes in 2D string theory, it is necessary to generalize what we have obtained above to the quantum theory. We have been succeeded in constructing a quantum version of the nonlinear $\hat{W}_\infty$ algebra, by quantizing the $SL(2, R)_k/U(1)$ coset model, with the bosonized currents

$$\begin{aligned} \psi_+(z; p) &= \frac{1}{2}[(1 + \sqrt{1 - 2p})\bar{j} + (1 - \sqrt{1 - 2p})j]e^{\sqrt{p}(\bar{\phi} + \phi)}, \\ \psi_-(z; p) &= \frac{1}{2}[(1 - \sqrt{1 - 2p})\bar{j} + (1 + \sqrt{1 - 2p})j]e^{-\sqrt{p}(\bar{\phi} + \phi)}. \end{aligned} \quad (30)$$

where $\phi(z)$ and $\bar{\phi}(z)$ are a pair of complex bosons, and their currents $\bar{j}(z)$, $j(z)$ satisfy

$$\bar{j}(z)j(z') \sim \frac{1}{(z - z')^2}, \quad \bar{j}(z)\bar{j}(z') \sim j(z)j(z') \sim 0. \quad (31)$$

Note the parameter $p \equiv k^{-1}$ plays the role of the Planck constant as the essential parameter in quantum corrections (or quantum deformation). We have included necessary quantum corrections in the currents. The classical limit is recovered by first rescaling $\phi \rightarrow \sqrt{k}\phi$, $\psi \rightarrow \sqrt{k}\psi$ and then letting $k \rightarrow \infty$.

Generalizing the classical equation (18), we use the whole OPE (up to all orders),

$$\psi_+(z; p)\psi_-(z'; p) = \epsilon^{-2p} \left\{ \epsilon^{-2} + \sum_{r=0}^{\infty} u_r(z; p) \frac{\epsilon^r}{r!} \right\} \quad (32)$$

(with $\epsilon \equiv z - z'$) to generate the quantum $\hat{W}_\infty(p)$ currents $u_r(z; p)$ in the KP basis. A closed expression for all $u_r(z; p)$, in terms of $\bar{j}$ and j only, has been derived in ref. 8.

The complete structure of the quantum $\hat{W}_\infty(p)$ algebra can be manifested by the OPE's between two currents $u_r(z; p)$ and $u_s(w; p)$, as usual in conformal field theory. In principle, these OPE's can be extracted from the OPE of four $SL(2, R)_k/U(1)$ currents:

$$(\epsilon\sigma)^{2p}(\psi_+(z; p)\psi_-(z - \epsilon; p))(\psi_+(w; p)\psi_-(w - \sigma; p)). \quad (33)$$

The closure of the quantum $\hat{W}_\infty(p)$ algebra is ensured by the closure of the OPE's associated with the enveloping algebra of the $SL(2, R)_k$ currents in the neutral sector in the conformal model. However, once we write the algebra in the form of OPE's among u_r and u_s , the currents u_r may be considered as independent of each other, and the associativity of the OPE's automatically leads to closed Jacobi identities. So our quantum version of the $\hat{W}_\infty$ algebra holds even beyond the context of the coset model.

The $p = 1$ case, corresponding to level $k = 1$, may be interpreted as the “typical” quantum case. For this value of p , a lot of expressions simplify.

Proposition 6: In the case of $p = 1$, we have

$$\begin{aligned} u_0(z)u_s(w) &= \frac{-1}{z-w}u'_s(w) + \text{terms in other powers of } (z-w)^{-1}, \\ u_1(z)u_s(w) &= \frac{-2}{z-w} \left[\sum_{l=1}^s (-1)^l \binom{s}{l} (u_0^{(l)} u_{s-l}) + \frac{u'_{s+1}}{(s+1)} + \frac{(-1)^s u_0^{(s+2)}}{(s+1)(s+2)} \right] (w) \\ &\quad + \text{terms in other powers of } (z-w)^{-1}. \end{aligned} \quad (34)$$

Commuting Quantum $\hat{W}_\infty$ Charges and Quantum KP Hierarchy

The above OPE's provide sufficient information for constructing an infinite set of commuting quantum $\hat{W}_\infty$ charges, whose densities are quantum deformation of the classical Hamiltonian functions (8). For generic p , it is not hard to construct

$$\begin{aligned} H_2(z; p) &= \frac{1}{(1-2p)} u_0(z; p), & H_3(z; p) &= u_1(z; p) + \frac{1}{2} u'_0(z; p), \\ H_4(z; p) &= u_2(z; p) + u'_1(z; p) + \frac{(5+4p)}{15} u''_0(z; p) + \frac{p}{(1-2p)} (u_0 u_0)(z; p). \end{aligned} \quad (35)$$

In the “typical” quantum case with $p = 1$, we can exploit the two general OPE (34) to explicitly construct an infinite number of commuting quantum $\hat{W}_\infty$ charges Q_m .

Let us recall that in conformal field theory the local product $(AB)(z)$ is defined by

$$(AB)(z) = \oint_z \frac{A(w)B(z)}{w-z} dw \quad (36)$$

in which the small contour of integration encircles z . This product is neither commutative nor associative. The symmetric local product of N local operators is defined to be the totally symmetrized sum of their multiple local products taken from the left:

$$\langle A_1 A_2 \cdots A_N \rangle = \frac{1}{N!} \sum_{P\{i\}} (\cdots ((A_{i_1} A_{i_2}) A_{i_3}) \cdots A_{i_N}) \quad (37)$$

where $P\{i\}$ denotes the summation over all possible permutations.

To construct quantum $\hat{W}_\infty$ charges, let us assign a degree 1 to ∂_z and $r+2$ to u_r , and assume that the quantum charge-density $H_m(z)$ is homogeneous and of degree m , and led by the highest-spin current u_{m-2} with unity coefficient. We require their charges commute with each other:

$$[\oint_0 H_n(z) dz, \oint_0 H_m(w) dw] = 0. \quad (38)$$

Proposition 7: The commutativity with Q_2 is always satisfied:

$$[\oint_0 H_2(z) dz, \oint_0 H_m(w) dw] = 0 \quad (39)$$

Thus the first set of nontrivial equations in eq.(38) start with $n = 3$. Here H_3 can only be u_1 plus a derivative of u_0 ; the latter does not contribute to the charge

Q_3 . We have shown⁸ that charges Q_m are completely determined by the requirement of commutativity with Q_3 alone. Both amusingly and amazingly, the so-determined charges Q_m automatically commute with each other. (A similar situation has happened before for commuting charges for the quantum KdV equation.¹⁰)

Proposition 8: The charges

$$Q_m = \oint_0 \sum_l \sum_{\{i\}} C_{i_1 i_2 \dots i_l}(m) \langle u_{i_1} u_{i_2} \dots u_{i_l} \rangle(z) dz \quad (40)$$

with the elegant expression for the coefficients

$$C_{i_1 i_2 \dots i_l}(m) = \frac{(-1)^{l-1} (l-1)! (m-2)!}{d_1! d_2! \dots d_k! i_1! i_2! \dots i_l!}, \quad (41)$$

provides a unique solution to eq.(38) for arbitrary m . Here for given number of currents in the product, l , the summation is over all partitions $\{i_k\}$ of $m-2l$, satisfying $i_1 + i_2 + \dots + i_l = m-2l$; and here d 's denote the degeneracies in the partition: $0 \leq i_1 = i_2 = \dots = i_{d_1} < i_{d_1+1} = \dots = i_{d_1+d_2} < \dots = i_{d_1+d_2+\dots+d_k(=l)}$.

We emphasize that in the proof of this Proposition, we have used only the OPE's in eq.(34); the two boson realization (32) of the $\hat{W}_\infty$ currents u_r is never used.

Now it is straightforward to construct a quantum version of the KP hierarchy in the Hamiltonian form (2). We use the quantum $\hat{W}_\infty$ as the Hamiltonian structure, and the densities of the quantum charges Q_m as corresponding Hamiltonian functions H_m . The latter naturally generate an infinite set of compatible flows in times t_m ($m = 1, 2, \dots$):

$$\frac{\partial u_r}{\partial t_m} = [u_r, Q_{m+1}]. \quad (42)$$

The mutual commutativity of these quantum charges implies that they are conserved charges of the flows (42):

$$\frac{\partial Q_n}{\partial t_m} = [Q_n, Q_{m+1}] = 0. \quad (43)$$

This ensures the complete integrability of the hierarchy (42). We call it the $p = 1$ quantum KP hierarchy and view it as a desired quantum version of the classical KP hierarchy. To justify this, we have computed the first three quantum KP charges for generic p and have shown that they coincide with the charges (40) at $p = 1$ and reduce

to the classical KP charges when $p \rightarrow 0$. Also we have calculated the first few quantum KP flows with generic p :

$$\begin{aligned}\frac{\partial u_0}{\partial t_1} &= (1-2p)u'_0, \quad \frac{\partial u_1}{\partial t_1} = (1-2p)u'_1, \quad \frac{\partial u_0}{\partial t_2} = (1-2p)(2u'_1 + u''_0), \\ \frac{\partial u_1}{\partial t_2} &= (2-3p)u'_2 + (1-p)u''_1 + \frac{(1+2p)p}{6}u'''_0 + p(u_0u_0)', \\ \frac{\partial u_0}{\partial t_3} &= 3(1-2p)(u'_2 + u''_1) + \frac{(3-4p-p^2)}{3}u'''_0 + 3p(u_0u_0)'.\end{aligned}\tag{44}$$

Manipulating the last three equations, we obtain the first dynamically nontrivial quantum evolution equation for the current $u_0(z; p)$:

$$6(2-3p)\frac{\partial u'_0}{\partial t_3} - 9\frac{\partial^2 u_0}{\partial t_2^2} = (3-p-16p^2+18p^3)u''''_0 + 18(1-p)p(u_0u_0)''.\tag{45}$$

Rescaling $u_0 \rightarrow u_0/p$ and taking $p \rightarrow 0$, this reduces to the classical KP equation, as claimed.

It should also be possible to obtain an alternative quantum deformation of the classical KP hierarchy through deforming its much simpler first Hamiltonian structure – the $W_{1+\infty}$. Because of its linearity, quantization of $W_{1+\infty}$ should be straightforward. In the KP basis, the complete structure of the quantum $W_{1+\infty}$ is neatly manifested by the following OPE's:

$$\begin{aligned}u_r(z)u_s(w) &= \sum_{l=0}^r \frac{r!}{(r-l)!} \frac{u_{r+s-l}(z)}{(z-w)^{l+1}} - \sum_{l=0}^s (-1)^l \frac{s!}{(s-l)!} \frac{u_{r+s-l}(w)}{(z-w)^{l+1}} \\ &\quad + \frac{(-1)^s cr!s!}{(z-w)^{r+s+2}} + O(z-w).\end{aligned}\tag{46}$$

The remaining issue is to construct an complete set of infinitely many commuting quantum charges in accordance to (46), in order for the associated quantum KP hierarchy to be integrable.

To conclude, we remark that it would be interesting to clarify the role of our world-sheet nonlinear $\hat{W}_\infty$ symmetry in physics of 2D black holes, and to invent a formalism for 2D strings in terms of the KP hierarchy or $\hat{W}_\infty$ constraints.

Acknowledgement: The work was supported in part by US NSF grant PHY-9008452.

REFERENCES

1. F. Yu and Y.-S. Wu, Phys. Lett. B263 (1991) 220.
2. F. Yu and Y.-S. Wu, Nucl. Phys. B373 (1992) 713.
3. F. Yu and Y.-S. Wu, Phys. Rev. Lett. 68 (1992) 2996.
4. F. Yu and Y.-S. Wu, Utah preprint UU-HEP-92/11, May 1992; to be published in Phys. Lett. B.
5. K. Yamagishi, Phys. Lett. B259 (1991) 436.
6. J. M. Figueroa-O'Farrill, J. Mas and E. Ramos, Phys. Lett. B266 (1991) 298; preprint BONN-HE-92/20, US-FT-92/7 or KUL-TF-92/20.
7. F. Yu and Y.-S. Wu, Utah preprint UU-HEP-92/07, August 1992.
8. F. Yu and Y.-S. Wu, Utah preprint UU-HEP-92/12, August 1992.
9. D. A. Depireux, Laval preprint LAVAL-PHY-21-92; J. M. Figueroa-O'Farrill, J. Mas and E. Ramos, preprint BONN-HE-92/17, US-FT-92/4 or KUL-TF-92/26.
10. R. Sasaki and I. Yamanaka, Commun. Math. Phys. 108 (1987) 691; Adv. Stud. in Pure Math. 16 (1988) 271.